

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q8. Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet | 62.4 | 2.65 | (56.9 - 67.5) | 1,017 | 1.7 |
| Q9. Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else | 12.9 | 1.30 | (10.5 - 15.7) | 1,493 | 1.5 |
| Q10. Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol | 23.2 | 1.50 | (20.3 - 26.3) | 1,492 | 1.4 |
| Q11. Percentage of students who drove a car or other vehicle one or more times during the past 30 days when they had been drinking alcohol | 8.5 | 0.99 | (6.7 - 10.7) | 1,480 | 1.4 |
| Q14. Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days | 8.8 | 1.00 | (7.0 - 11.0) | 1,493 | 1.4 |
| Q15. Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school | 4.5 | 0.77 | (3.2 - 6.3) | 1,493 | 1.4 |
| Q17. Percentage of students who were in a physical fight one or more times during the past 12 months | 25.9 | 1.59 | (22.8 - 29.2) | 1,482 | 1.4 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|---|------------|----------------|-------------------------|--------------------|----------------|
| Q18. Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse | 3.9 | 0.51 | (3.0 - 5.1) | 1,487 | 1.0 |
| Q19. Percentage of students who were in a physical fight on school property one or more times during the past 12 months | 9.1 | 0.87 | (7.5 - 11.0) | 1,486 | 1.2 |
| Q20. Percentage of students who were ever hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend during the past 12 months | 9.6 | 0.89 | (8.0 - 11.6) | 1,490 | 1.2 |
| Q21. Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to | 7.0 | 0.80 | (5.6 - 8.8) | 1,483 | 1.2 |
| Q22. Percentage of students who had ever been bullied on school property during the past 12 months | 22.1 | 1.53 | (19.2 - 25.3) | 1,489 | 1.4 |
| Q23. Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months | 25.1 | 1.19 | (22.8 - 27.6) | 1,491 | 1.1 |
| Q24. Percentage of students who seriously considered attempting suicide during the past 12 months | 12.1 | 1.09 | (10.0 - 14.4) | 1,482 | 1.3 |
| Q25. Percentage of students who made a plan about how they would attempt suicide during the past 12 months | 9.8 | 1.07 | (7.8 - 12.2) | 1,490 | 1.4 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|---|------------|----------------|-------------------------|--------------------|----------------|
| Q26. Percentage of students who actually attempted suicide one or more times during the past 12 months | 4.7 | 0.83 | (3.3 - 6.6) | 1,412 | 1.5 |
| Q27. Percentage of students who made a suicide attempt during the past 12 months that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse | 1.6 | 0.44 | (1.0 - 2.8) | 1,407 | 1.3 |
| Q29. Percentage of students who smoked a whole cigarette for the first time before age 13 years | 10.4 | 0.95 | (8.7 - 12.5) | 1,483 | 1.2 |
| Q30. Percentage of students who smoked cigarettes on one or more of the past 30 days | 20.8 | 1.56 | (17.8 - 24.1) | 1,488 | 1.5 |
| Q36. Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days | 8.4 | 1.02 | (6.6 - 10.7) | 1,490 | 1.4 |
| Q38. Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days | 16.1 | 1.19 | (13.9 - 18.7) | 1,490 | 1.3 |
| Q39. Percentage of students who had at least one drink of alcohol on one or more days during their life | 68.5 | 1.81 | (64.8 - 72.0) | 1,472 | 1.5 |
| Q40. Percentage of students who had their first drink of alcohol other than a few sips before age 13 years | 14.8 | 0.98 | (12.9 - 16.8) | 1,483 | 1.1 |
| Q41. Percentage of students who had at least one drink of alcohol on one or more of the past 30 days | 39.3 | 2.18 | (35.0 - 43.8) | 1,392 | 1.7 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q42. Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days | 24.0 | 1.68 | (20.8 - 27.5) | 1,469 | 1.5 |
| Q43. Among students who reported current alcohol use, the percentage who usually got the alcohol they drank from someone who gave it to them during the past 30 days | 31.1 | 2.14 | (26.9 - 35.5) | 518 | 1.1 |
| Q44. Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days | 4.3 | 0.68 | (3.2 - 5.9) | 1,484 | 1.3 |
| Q45. Percentage of students who used marijuana one or more times during their life | 40.5 | 2.02 | (36.6 - 44.6) | 1,471 | 1.6 |
| Q46. Percentage of students who tried marijuana for the first time before age 13 years | 8.4 | 0.82 | (6.9 - 10.2) | 1,468 | 1.1 |
| Q47. Percentage of students who used marijuana one or more times during the past 30 days | 25.6 | 1.86 | (22.1 - 29.5) | 1,476 | 1.6 |
| Q48. Percentage of students who used marijuana on school property one or more times during the past 30 days | 6.8 | 0.78 | (5.4 - 8.6) | 1,476 | 1.2 |
| Q49. Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life | 6.5 | 0.79 | (5.1 - 8.3) | 1,482 | 1.2 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q50. Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during the past 30 days | 4.0 | 0.71 | (2.8 - 5.6) | 1,483 | 1.4 |
| Q51. Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life | 11.9 | 1.19 | (9.7 - 14.5) | 1,490 | 1.4 |
| Q52. Percentage of students who used heroin one or more times during their life | 3.0 | 0.54 | (2.1 - 4.3) | 1,491 | 1.2 |
| Q53. Percentage of students who used methamphetamines one or more times during their life | 4.7 | 0.60 | (3.6 - 6.0) | 1,491 | 1.1 |
| Q54. Percentage of students who used ecstasy one or more times during their life | 6.8 | 0.99 | (5.1 - 9.1) | 1,489 | 1.5 |
| Q55. Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life | 2.1 | 0.40 | (1.4 - 3.1) | 1,491 | 1.1 |
| Q57. Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months | 22.1 | 1.44 | (19.4 - 25.1) | 1,480 | 1.3 |
| Q58. Percentage of students who ever had sexual intercourse | 46.3 | 2.10 | (42.1 - 50.5) | 1,444 | 1.6 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q59. Percentage of students who had sexual intercourse for the first time before age 13 years | 4.3 | 0.65 | (3.2 - 5.8) | 1,439 | 1.2 |
| Q60. Percentage of students who had sexual intercourse with four or more people during their life | 11.4 | 1.17 | (9.3 - 14.0) | 1,443 | 1.4 |
| Q61. Percentage of students who had sexual intercourse with one or more people during the past three months | 36.3 | 1.97 | (32.4 - 40.3) | 1,442 | 1.6 |
| Q62. Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse | 21.4 | 1.77 | (18.1 - 25.1) | 520 | 1.0 |
| Q63. Among students who had sexual intercourse during the past three months, the percentage who used a condom during last sexual intercourse | 56.8 | 2.59 | (51.6 - 61.9) | 513 | 1.2 |
| Q64. Among students who had sexual intercourse during the past three months, the percentage who used birth control pills to prevent pregnancy before last sexual intercourse | 33.0 | 2.80 | (27.6 - 38.8) | 492 | 1.3 |
| Q65. Percentage of students who described themselves as slightly or very overweight | 28.4 | 1.37 | (25.7 - 31.2) | 1,485 | 1.2 |
| Q66. Percentage of students who were trying to lose weight | 44.0 | 1.29 | (41.5 - 46.6) | 1,485 | 1.0 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q67. Percentage of students who exercised to lose weight or to keep from gaining weight during the past 30 days | 59.9 | 1.37 | (57.2 - 62.7) | 1,486 | 1.1 |
| Q68. Percentage of students who ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight during the past 30 days | 40.6 | 1.09 | (38.4 - 42.8) | 1,486 | 0.9 |
| Q69. Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight during the past 30 days | 8.8 | 0.89 | (7.2 - 10.7) | 1,473 | 1.2 |
| Q70. Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days | 5.1 | 0.66 | (3.9 - 6.6) | 1,471 | 1.1 |
| Q71. Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days | 3.9 | 0.77 | (2.7 - 5.8) | 1,473 | 1.5 |
| Q78. Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days | 22.1 | 1.24 | (19.8 - 24.7) | 1,488 | 1.1 |
| Q79. Percentage of students who drank three or more glasses per day of milk during the past seven days | 21.3 | 1.33 | (18.7 - 24.1) | 1,486 | 1.3 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q80. Percentage of students who were physically active for a total of at least 60 minutes per day on five or more of the past seven days | 45.3 | 2.10 | (41.1 - 49.5) | 1,488 | 1.6 |
| Q81. Percentage of students who watched three or more hours per day of TV on an average school day | 23.0 | 1.32 | (20.5 - 25.7) | 1,483 | 1.2 |
| Q82. Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day | 23.9 | 1.12 | (21.8 - 26.3) | 1,486 | 1.0 |
| Q83. Percentage of students who attended physical education (PE) classes on one or more days in an average week when they were in school | 41.1 | 3.32 | (34.7 - 47.9) | 1,478 | 2.6 |
| Q85. Percentage of students who had ever been taught in school about AIDS or HIV infection | 89.9 | 1.07 | (87.5 - 91.9) | 1,480 | 1.4 |
| Q88. Percentage of students who would describe their grades in school as mostly D's and F's during the past 12 months | 4.7 | 0.64 | (3.6 - 6.2) | 1,480 | 1.2 |
| Q89. Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life | 20.4 | 1.29 | (17.9 - 23.1) | 1,478 | 1.2 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q90. Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during the past 30 days | 10.4 | 1.24 | (8.1 - 13.1) | 1,477 | 1.6 |
| Q91. Percentage of students who have taken over-the-counter drugs to get high one or more times during their life | 11.9 | 1.22 | (9.6 - 14.5) | 1,484 | 1.5 |
| Q92. Percentage of students who have taken over-the-counter drugs to get high one or more times during the past 30 days | 6.1 | 0.86 | (4.5 - 8.0) | 1,485 | 1.4 |
| Q93. Percentage of students who have had sexual contact with females and males during their life | 3.8 | 0.62 | (2.8 - 5.3) | 1,471 | 1.2 |
| Q94. Percentage of students who most of the time or always went hungry during the past 30 days because there was not enough food in their home | 4.1 | 0.64 | (3.0 - 5.6) | 1,470 | 1.2 |
| Q95. Percentage of students who last saw a doctor or nurse for a check-up or physical exam when they were not sick or injured during the past 12 months | 68.0 | 1.80 | (64.3 - 71.5) | 1,479 | 1.5 |
| Q96. Percentage of students who last saw a dentist for a check-up, exam, teeth cleaning, or other dental work during the past 12 months | 82.4 | 1.38 | (79.5 - 85.0) | 1,476 | 1.4 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|--|------------|----------------|-------------------------|--------------------|----------------|
| Q97. Percentage of students who strongly agree or agree that their parents or other adults in their family have clear rules and standards for their behavior | 77.2 | 1.84 | (73.3 - 80.6) | 1,475 | 1.7 |
| Q98. Percentage of students who performed any organized community service as a non-paid volunteer, such as serving meals to the elderly, picking up litter, helping out at a hospital, or building homes for the poor, one or more times during the past 30 days | 36.4 | 1.88 | (32.7 - 40.3) | 1,472 | 1.5 |
| Q99. Percentage of students who participate in one or more clubs or organizations outside of school (other than sports teams), such as 4-H, Boys and Girls Club, YMCA, or, church groups, during an average week when they are in school | 38.4 | 1.98 | (34.5 - 42.4) | 1,475 | 1.6 |
| Q100. Percentage of students who strongly agree or agree that in their community they feel like they matter to people | 41.5 | 1.81 | (37.9 - 45.1) | 1,472 | 1.4 |
| Q101. Percentage of students who think people are at no risk of harming themselves (physically or in other ways) if they smoke one or more packs of cigarettes per day | 4.2 | 0.65 | (3.1 - 5.7) | 1,473 | 1.2 |
| Q102. Percentage of students who think people are at no risk of harming themselves (physically or in other ways) if they smoke marijuana occasionally | 23.1 | 1.66 | (19.9 - 26.6) | 1,470 | 1.5 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|---|------------|----------------|-------------------------|--------------------|----------------|
| Q103. Percentage of students who think people are at no risk of harming themselves (physically or in other ways) if they smoke marijuana regularly | 12.9 | 1.26 | (10.6 - 15.6) | 1,475 | 1.4 |
| Q104. Percentage of students who think people are at no risk of harming themselves (physically or in other ways) if they have one or two drinks of an alcoholic beverage (beer, wine, or liquor) each weekend | 27.3 | 1.53 | (24.3 - 30.5) | 1,474 | 1.3 |
| Q105. Percentage of students who think people are at no risk of harming themselves (physically or in other ways) if they have one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day | 7.6 | 0.98 | (5.8 - 9.8) | 1,468 | 1.4 |
| Q106. Percentage of students who think people are at no risk of harming themselves (physically or in other ways) if they have five or more drinks of an alcoholic beverage (beer, wine, or liquor) each weekend | 8.2 | 0.84 | (6.7 - 10.1) | 1,466 | 1.2 |
| Q107. Percentage of students who think people are at no risk of harming themselves (physically or in other ways) if they have five or more drinks of an alcoholic beverage (beer, wine, or liquor) nearly every day | 4.7 | 0.82 | (3.3 - 6.7) | 1,466 | 1.5 |
| Q108. Percentage of students who think it is very wrong or wrong for someone their age to drink beer, wine, or liquor regularly | 48.8 | 1.78 | (45.3 - 52.4) | 1,471 | 1.4 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|---|------------|----------------|-------------------------|--------------------|----------------|
| Q109. Percentage of students who think it is very wrong or wrong for someone their age to smoke cigarettes | 66.4 | 2.10 | (62.0 - 70.4) | 1,468 | 1.7 |
| Q110. Percentage of students who think it is very wrong or wrong for someone their age to smoke marijuana | 57.1 | 1.75 | (53.5 - 60.5) | 1,467 | 1.4 |
| Q111. Percentage of students who think it is very wrong or wrong for someone their age to use LSD, cocaine, amphetamines, or another illegal drug | 88.5 | 1.29 | (85.7 - 90.9) | 1,469 | 1.6 |
| Q112. Percentage of students whose parents think it is very wrong or wrong for someone their age to drink beer, wine, or liquor regularly | 82.4 | 1.33 | (79.5 - 84.9) | 1,467 | 1.3 |
| Q113. Percentage of students whose parents think it is very wrong or wrong for someone their age to smoke cigarettes | 85.7 | 1.42 | (82.6 - 88.3) | 1,468 | 1.6 |
| Q114. Percentage of students whose parents think it is very wrong or wrong for someone their age to smoke marijuana | 84.7 | 1.07 | (82.5 - 86.7) | 1,471 | 1.1 |
| Q115. Percentage of students who think it would be very easy for them to get some beer, wine, or liquor if they wanted to | 37.9 | 1.58 | (34.8 - 41.1) | 1,467 | 1.2 |
| Q116. Percentage of students who think it would be very easy for them to get some cigarettes if they wanted to | 49.5 | 2.40 | (44.8 - 54.3) | 1,469 | 1.8 |

*The average design effect across all items for this survey is 1.4.

2009 YOUTH RISK BEHAVIOR SURVEY RESULTS

New Hampshire High School Survey

Sample Statistics

| Question | Percentage | Standard Error | 95% Confidence Interval | Number of Students | Design Effect* |
|---|------------|----------------|-------------------------|--------------------|----------------|
| Q117. Percentage of students who think it would be very easy for them to get some marijuana if they wanted to | 43.8 | 1.73 | (40.3 - 47.3) | 1,469 | 1.3 |
| Q118. Percentage of students who think it would be very easy for them to get some drug like cocaine, LSD, or amphetamines if they wanted to | 10.4 | 1.25 | (8.2 - 13.2) | 1,465 | 1.6 |

*The average design effect across all items for this survey is 1.4.